# NARCAN (NALOXONE)

**OPIOID OVERDOSE** 




### WHAT IS FENTANYL



- FENTANYL, HEROIN'S SYNTHETIC COUSIN, IS 50-100 TIMES MORE POTENT THAN MORPHINE
- HEROIN IS BEING SOLD LACED WITH FENTANYL
- FENTANYL IS BEING SOLD UNDER THE GUISE THAT IT IS HEROIN
- LAWS PASSED A FEW YEARS AGO TO ADDRESS MISUSE OF PRESCRIPTION DRUGS

# HOW NARCAN WORKS



# NARCAN 4MG DOSES

PERSONNEL WILL BE ISSUED TWO 4MG DOSES OF NARCAN
THIS SPRAY DISPENSES A FINE MIST INTO THE NASAL PASSAGEWAY
WHICH IS THEN ABSORBED DIRECTLY INTO THE BLOODSTREAM

KEEP THE TWO DOSES TOGETHER AND KEEP OUT OF DIRECT SUNLIGHT OR HIGH HEAT FOR EXTENDED PERIODS BECAUSE IT CAN REDUCE THE EFFECTIVENESS OF THE DOSE



# INDICATIONS OF OPIOID O.D.

PATIENT IS UNRESPONSIVE WITH MINIMAL OR NO RESPIRATIONS MAY OR MAY NOT HAVE A PULSE/ CONSTRICTED PUPILS LIPS MAY BE BLUE OR GREY AND PATIENT HAS POOR COLOR PILL BOTTLES, NEEDLES, FAMILY/FRIENDS INDICATE OPIOID USE



**SCENE SAFETY!!** 

PPE IS ESSENTIAL

START LIFE SAVING MEASURES

**ENSURE RESCUE IS RESPONDING** 

#### NARCAN USE IS INDICATED

**ADMINISTER FULL 4 MG DOSE** 

**IT TAKES 2-5 MINUTES TO BE EFFECTIVE** 

IF NORMAL BREATHING IS NOT RESTORED AND PARAMEDICS ARE NOT ON SCENE, A SECOND DOSE MAY BE GIVEN



# PROPER US OF NARCAN






## PROPER USE OF NARCAN SPRAY

DO NOT TEST OR PRIME THE NARCAN AS IT WILL WASTE THE DOSE SPRAY IN EITHER NOSTRIL AND ONCE FULL DOSE HAS BEEN GIVEN CONTINUE WITH RESCUE BREATHING AND OR CPR IF INDICATED \*\*NOT FOR USE OF INFANTS (12 MONTHS AND YOUNGER)

IF PATIENT IS BREATHING, PUT THEM IN THE RECOVERY POSITION

# THE EFFECTS OF NARCAN CAN WEAR OFF SO CONTINUE TO MONITOR THE PATIENT AND IF BREATHING, PUT IN RECOVERY POSITION



# **DOCUMENTATION**

IF NARCAN IS USED- COMPLETE THE "NARCAN USE REPORT" SH-R-630 AND FORWARD IT TO THE DEPARTMENT NARCAN COORDINATOR AT SEB/ESD

IF A SH-AD 49 IS WRITTEN, YOUR USE OF NARCAN SHOULD BE DOCUMENTED IN THE REPORT REQUEST REPLACEMENT NARCAN AND/OR CPR KITS THROUGH THE DEPARTMENT NARCAN COORDINATOR