## Additional and Booster Doses for Fully Vaccinated Persons Who Received a Non-FDA Authorized/Approved Vaccine for Their Primary Series

	WHO-listed COVID-19 vaccine <sup>2</sup>	Active vaccine from a COVID-19 clinical trial <sup>3</sup>
Additional (3 <sup>rd</sup> ) Dose <sup>4</sup>	Who: Strongly recommended if you are age 12 or older and have a moderately or severely weakened immune system.	
	When: At least 28 days after your second dose	
Booster Dose <sup>4</sup>	<ul> <li>Who: Strongly Recommended for everyone age 18 and older.</li> <li>Very important if you are:         <ul> <li>Age 65 and older</li> <li>Age 18 and older and have an underlying medical condition or have risk of exposure at work or in the community.</li> </ul> </li> <li>When: At least 6 months after your second dose<sup>5</sup>.</li> </ul>	

- 1 You are considered fully vaccinated two weeks after receiving your final dose of the vaccine series. See When You've Been Fully Vaccinated webpage for more details.
- This also includes if you completed a mix and match series. A mix and match series is a combination of FDA-approved, FDA-authorized, or WHO-listed COVID-19 vaccines. Current non-FDA approved/authorized WHO-listed vaccines are AstraZeneca-Oxford (e.g., Covishield, Vaxzevria), Sinopharm, Sinovac, and Covaxin. See the When Am I Fully <u>Vaccinated?</u> section on the When You've Been Fully Vaccinated webpage to learn more.
- 3 If you took part in the AstraZeneca or Novavax clinical vaccine trial and you received the "active" COVID-19 vaccine (not placebo).
- The Pfizer COVID-19 vaccine is the only vaccine authorized as an additional dose or booster for people who did not receive an FDA authorized/approved COVID-19 vaccine series.
- If you are immunocompromised, this will be 6 months after your additional (3<sup>rd</sup>) dose.

For more information, see How to get Vaccinated, Doses for Immunocompromised and Booster Doses.

