

## **COVID-19 VACCINE ELIGIBILITY**

## Primary, Additional and Booster Doses for FDA Approved/Authorized COVID-19 Vaccines<sup>1</sup>

	Pfizer <sup>2</sup> /Comirnaty	<b>Moderna</b> <sup>3</sup>	Johnson & Johnson (J&J)
	Preferred vaccines for primary series, additional doses, and booster doses.		For people who cannot get a Pfizer or Moderna vaccine. <sup>4</sup>
Eligible age	5 and older <sup>2</sup>	18 and older	18 and older
Primary Series	Two doses 21 days apart	Two doses 28 days apart	One dose
Booster Dose	<ul> <li>Who: Everyone age 12 and older who got Pfizer for their primary series should get a booster dose.</li> <li>People 18+ can get any of the 3 vaccines: Pfizer or Moderna are preferred over J&amp;J booster.</li> <li>People age 12-17 can get the Pfizer vaccine.</li> </ul>	Who: Everyone who got Moderna for their primary series should get a booster dose.  Can get any of the 3 vaccines: Pfizer or Moderna are preferred over J&J booster	Who: Everyone who got J&J for their primary series should get a booster dose.  Can get any of the 3 vaccines: Pfizer or Moderna are preferred over J&J booster  This includes if you have a moderately or severely weakened immune system
	<b>When:</b> 5 months after your second dose <sup>6</sup>	<i>When:</i> 5 months after your second dose <sup>6</sup>	<b>When</b> : 2 months after your J&J dose
Additional (3 <sup>rd</sup> ) Dose for people with weak immune systems <sup>3,5</sup>	Who: People age 5 or older who have a moderately or severely weakened immune system should get an additional dose of the same vaccine as their primary series. Children 5-17 should only get Pfizer vaccine. <sup>2</sup> When: At least 28 days after your second dose		Not recommended. See Booster Dose above

- 1 If you received a WHO-listed vaccine or a vaccine as part of a COVID-19 vaccine clinical trial, see table on the next page for additional dose and booster dose information.
- The Pfizer vaccine for children 5-11 years of age has the same active ingredients as the adult vaccine but is a smaller dose  $(1/3^{rd})$  the dose that teens and adults receive).
- **3** Moderna doses vary. The primary series and the additional (3<sup>rd</sup>) dose are a full dose. The booster dose is a half dose.
- 4 The J&J vaccine is available for people who can't get a Pfizer or Moderna vaccine for medical reasons (such as severe allergic reaction to a vaccine ingredient) and for people who prefer it. But the Pfizer and Moderna COVID-19 vaccines are recommended for both primary and booster vaccination. This is because they offer better protection against COVID-19 than the J&J vaccine and the risks are lower. Getting any COVID-19 vaccine is better than remaining unvaccinated.
- **Additional dose** The CDC recommends that moderately to severely immunocompromised people who received an mRNA vaccine (Pfizer or Moderna) as their primary series receive an additional (3<sup>rd</sup>) dose. The additional dose should be the same COVID-19 vaccine as their primary series. Talk to your doctor about the need to get an additional dose of COVID-19 vaccine and ask about the best timing based on your current treatment plan. *Note*: if you are immunocompromised and received the J&J vaccine for your primary series, you should get a booster dose. A Pfizer or Moderna booster dose is preferred.
- 6 If you are immunocompromised, this will be after your additional (3<sup>rd</sup>) dose.

For more information, see How to get Vaccinated, Doses for Immunocompromised and Booster Doses.

