



COVID-19 VACCINE ELIGIBILITY



Additional and Booster Doses for Fully Vaccinated¹ Persons Who Received a Non-FDA Authorized/Approved Vaccine for Their Primary Series

	WHO-listed COVID-19 vaccine ²	Active vaccine from a COVID-19 clinical trial ³
Additional (3rd) Dose ⁴	Who: People age 12 or older and have a moderately or severely weakened immune system should get a 3 rd dose	
	When: At least 28 days after your second dose	
Booster Dose ⁴	Who: People age 12 and older should get a booster dose.	
	When: 5 months after your second dose ⁵	

- 1 You are considered fully vaccinated** two weeks after receiving your final dose of the vaccine series. See [When You've Been Fully Vaccinated](#) webpage for more details.
- 2 This also includes if you completed a mix and match series** – this is a combination of FDA-approved, FDA-authorized, or WHO-listed COVID-19 vaccines. Current non-FDA approved/authorized [WHO-listed vaccines](#) include AstraZeneca-Oxford (Vaxzevria), Serum Institute of India Pvt. Ltd (Covishield and Covovax), BIBP (Sinopharm), Sinovac (CoronaVac), Bharat Biotech International (Covaxin), and Novavax (Nuvaxovid). See the [When Am I Fully Vaccinated?](#) section on the *When You've Been Fully Vaccinated* webpage to learn more.
- 3 If you took part in a clinical vaccine trial** and you received all of the recommended “active” COVID-19 vaccine (not placebo). Note: the vaccine should be either a WHO-listed vaccine that is not FDA-approved/authorized or a vaccine for which a U.S. data and safety monitoring board or equivalent has independently confirmed efficacy.
- 4 The Pfizer COVID-19 vaccine is the only vaccine authorized** as an additional dose or booster for people who did not receive an FDA authorized/approved COVID-19 vaccine series.
- 5 If you are immunocompromised**, this will be 5 months after your additional (3rd) dose.

For more information, see [How to get Vaccinated](#), [Doses for Immunocompromised](#) and [Booster Doses](#).

